arXiv:1010.5728v1 [math.DG] 27 Oct 2010

ON AFFINE CONNECTIONS IN A RIEMANNIAN MANIFOLD WITH A CIRCULANT METRIC AND TWO CIRCULANT AFFINOR STRUCTURES[1]

IVA DOKUZOVA, DIMITAR RAZPOPOV

Abstract. In the present paper it is considered a class V of 3-dimensional Riemannian manifolds M with a metric g and two affinor tensors q and S. It is defined another metric $\bar{g}$ in M. The local coordinates of all these tensors are circulant matrices. It is found:

1) a relation between curvature tensors R and $\bar{R}$ of g and $\bar{g}$, respectively;

2) an identity of the curvature tensor R of g in the case when the curvature tensor $\bar{R}$ vanishes;

3) a relation between the sectional curvature of a 2-section of the type $\{x, qx\}$ and the scalar curvature of M.

1. Preliminaries

We consider a 3-dimensional Riemannian manifold M with a metric tensor g and two affinor tensors q and S such that: their local coordinates form circulant matrices. So these matrices are as follows:

$$g_{ij} = \begin{pmatrix} A & B & B \\ B & A & B \\ B & B & A \end{pmatrix}, \quad A > B > 0, \tag{1}$$

where A and B are smooth functions of a point $p(x^1, x^2, x^3)$ in some $F \subset R^3$,

$$q_{i}^{\cdot j} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \qquad S_{i}^{j} = \begin{pmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{pmatrix}. \tag{2}$$

Let ∇ be the connection of g. The following results have been obtained in [1].

$$q^3 = E; \quad g(qu, qv) = g(u, v), \quad u,\ v \in \chi M. \tag{3}$$

$$\nabla q = 0 \quad \Leftrightarrow \quad gradA = gradB.S. \tag{4}$$

$$0 < B < A \quad \Rightarrow \quad g\ is\ possitively\ defined. \tag{5}$$

---

[1] This work is partially supported by project RS09 - FMI - 003 of the Scientific Research Fund, Paisii Hilendarski University of Plovdiv, Bulgaria

If M has a metric g from (1), affinor structures q and S from (2) and $\nabla q = 0$, then we note for brevity, that M is in the class V.

We denote $\tilde{q}^s_j = q^s_a q^a_j$, $\Phi^s_j = q^s_j + \tilde{q}^s_j$, and from (2) we have:

$$\tilde{q}^s_j = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \qquad \Phi^s_j = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}. \tag{6}$$

2. On affine connections in a Riemannian manifold

Let M be in V. We note $f_{ij} = g_{ik}q^k_j + g_{jk}q^k_i$, i.e.

$$f_{ij} = \begin{pmatrix} 2B & A+B & A+B \\ A+B & 2B & A+B \\ A+B & A+B & 2B \end{pmatrix}. \tag{7}$$

We calculate $\det f_{ij} = 2(A-B)^2(A+2B) \neq 0$, so f is non-degenerated symmetric tensor field. Evidently, we have that $\nabla q = 0$, which thank's to (2), (6), (7), implies:

$$\nabla \tilde{q} = 0, \qquad \nabla f = 0, \qquad \nabla S = 0, \qquad \nabla \Phi = 0. \tag{8}$$

For later use, from (1) – (7), we find the next identities:

$$\Phi^s_j g_{is} = f_{ji}, \qquad \Phi^s_j f_{is} = 2g_{ji} + f_{ji}, \qquad f_{ji}g^{is} = \Phi^s_j \qquad g_{ji}f^{is} = \frac{1}{2}S^s_j. \tag{9}$$

Further, we suppose α and β are two smooth functions in F. Now, we construct the metric $\bar{g}$, as follows

$$\bar{g} = \alpha.g + \beta.f. \tag{10}$$

The local coordinates of $\bar{g}$ are

$$\bar{g}_{ij} = \begin{pmatrix} \alpha A + 2\beta B & \beta A + (\alpha+\beta)B & \beta A + (\alpha+\beta)B \\ \beta A + (\alpha+\beta)B & \alpha A + 2\beta B & \beta A + (\alpha+\beta)B \\ \beta A + (\alpha+\beta)B & \beta A + (\alpha+\beta)B & \alpha A + 2\beta B \end{pmatrix}.$$

In [2] it is proved the next assertion.

Theorem 2.1. *Let M be a manifold in V, also g and $\bar{g}$ be two metrics of M, related by (10). Let ∇ and $\overline{\nabla}$ be the corresponding connections of g and $\bar{g}$. Then $\overline{\nabla} q = 0$ if and only if, when*

$$grad\alpha = grad\beta.S. \tag{11}$$

Let $\beta = 0$ in (10). Then we have

$$\bar{g} = \alpha.g. \tag{12}$$

The condition (12) defines the well known conformal transformation in the Riemannian manifold M.

So, we will consider the case $\alpha = 0$ in (10). We obtain

$$\bar{g}_{ij} = \beta . f_{ij}. \tag{13}$$

Then from (11) we can get that $\overline{\nabla} q = 0$ if and only if, when β is a constant. Further, we are interested in the case $\overline{\nabla} q \neq 0$, i.e. β isn't a constant. Thank's to (13) we get

$$\nabla_k \bar{g}_{ij} = \beta_k f_{ij}, \qquad \beta_k = \nabla_k \beta. \tag{14}$$

We have the well known identities:

$$\overline{\nabla}_k \bar{g}_{ij} = \partial_k \bar{g}_{ij} - \overline{\Gamma}^a_{ki} \bar{g}_{aj} - \overline{\Gamma}^a_{kj} \bar{g}_{ai}, \tag{15}$$

$$\nabla_k \bar{g}_{ij} = \partial_k \bar{g}_{ij} - \Gamma^a_{ki} \bar{g}_{aj} - \Gamma^a_{kj} \bar{g}_{ai}, \tag{16}$$

$$\overline{\nabla}_k \bar{g}_{ij} = 0. \tag{17}$$

Using (9), (14) – (17), for the tensor $T^s_{ik} = \overline{\Gamma}^s_{ki} - \Gamma^s_{ki}$ of the affine deformation of ∇ and $\overline{\nabla}$, we find

$$T^s_{ik} = \beta_k \delta^s_i + \beta_i \delta^s_k - \frac{1}{2} \beta^a S^s_a f_{ik}, \qquad \beta_k \sim \frac{\beta_i}{2\beta}. \tag{18}$$

We have that $\overline{\nabla}_i q^k_j = \nabla_i q^k_j - \beta_j q^k_i + \tilde{\beta}_j \delta^k_i - \dfrac{1}{2} \beta^a S^k_a q^t_j f_{ti} + \dfrac{1}{2} \beta^a S^t_a q^k_t f_{ij}$.

Let R be the curvature tensor field of ∇. Let $\bar{R}$ be the curvature tensor field of $\overline{\nabla}$. It is well known the relation (see [3])

$$\overline{R}^h_{ijk} = R^h_{ijk} + \nabla_j T^h_{ik} - \nabla_k T^h_{ij} + T^s_{ik} T^h_{sj} - T^s_{ij} T^h_{sk}. \tag{19}$$

From (18) and (19) after some calculations we obtain

$$\begin{aligned} \overline{R}^h_{ijk} &= R^h_{ijk} + \delta^h_k (\nabla_j \beta_i - \beta_i \beta_j + \varphi f_{ij}) - \delta^h_j (\nabla_k \beta_i - \beta_i \beta_k + \varphi f_{ik}) \\ &\quad + \frac{1}{2} f_{ij} S^h_t (\nabla_k \beta^t - \beta_k \beta^t) - \frac{1}{2} f_{ik} S^h_t (\nabla_j \beta^t - \beta_j \beta^t), \quad \varphi = \frac{1}{2} \beta^t \beta_s S^s_t. \end{aligned} \tag{20}$$

Theorem 2.2. *Let M be in V, also ∇ and $\overline{\nabla}$ be the Riemannian connections of g and $\bar{g}$, related by (13). If $\overline{\nabla}$ is a locally flat connection, then curvature tensor field R of ∇ is*

$$\begin{aligned} R(x,y,z,u) = \frac{\tau}{6} [&(2g(x,u) g(y,z) - 2g(x,z) g(y,u) \\ &+ (g(qx,u) + g(x,qu))(g(qy,z) + g(y,qz)) \\ &- (g(qx,z) + g(x,qz))(g(qy,u) + g(y,qu))]. \end{aligned} \tag{21}$$

Proof. We have $\overline{R} = 0$. From (20) we find

$$R^h_{ijk} = \delta^h_j P_{ki} - \delta^h_k P_{ij} - f_{ij} Q^h_k + f_{ik} Q^h_j, \tag{22}$$

where $P_{ki} = \nabla_k \beta_i - \beta_i \beta_k + \varphi f_{ik}$, $Q^h_k = \dfrac{1}{2} S^h_t (\nabla_k \beta^t - \beta_k \beta^t)$.

Now, we contract $k = h$ in (22) and with the help of (9) we get

$$R_{ij} = -P_{ij} - \psi f_{ij}, \qquad \psi = \frac{1}{2} S_t^h \nabla_h \beta^t. \tag{23}$$

We note that $R_{ij} = R_{ijk}^k$ are the local coordinates of the Ricci tensor of ∇, also $\tau = R_{ij} g^{ij}$ and $\tau^* = R_{ij} f^{ij}$ are the first and the second scalar curvatures of M, respectively. The identity (23) implies

$$\tau^* = -2\varphi - 4\psi. \tag{24}$$

Using (9) we have that $Q_k^h = P_{ka} f^{ah} - \varphi \delta_k^h$, and from (23) we get

$$Q_k^h = -R_{ka} f^{ah} - (\psi + \varphi)\delta_k^h. \tag{25}$$

We substitute (23) – (25) in (22), and we find

$$R_{ijk}^h = \delta_k^h (R_{ij} - \frac{\tau^*}{2} f_{ij}) - \delta_j^h (R_{ki} - \frac{\tau^*}{2} f_{ki}) + f_{ij} R_{ka} f^{ah} - f_{ik} R_{ja} f^{ah}. \tag{26}$$

From (26) and $R_k^h = R_{ijk}^h g^{ij}$ we have

$$2R_k^h = \tau \delta_k^h + \frac{\tau^*}{2} \Phi_k^h - \Phi_k^t R_{ta} f^{ah}. \tag{27}$$

Now, we contract (27) with f_{ih}, and from identity $f_{ih} R_k^h = \Phi_i^a R_{ka}$ we obtain:

$$2\Phi_i^a R_{ka} = (\frac{\tau^*}{2} + \tau) f_{ki} + \tau^* g_{ki} - \Phi_k^t R_{ti}$$

and

$$2\Phi_k^a R_{ia} = (\frac{\tau^*}{2} + \tau) f_{ki} + \tau^* g_{ki} - \Phi_i^t R_{tk}.$$

The last system of two equations implies

$$\Phi_k^a R_{ia} = \frac{1}{3}((\frac{\tau^*}{2} + \tau) f_{ki} + \tau^* g_{ki}). \tag{28}$$

From (9) and (28) we find

$$\Phi_k^a R_{ia} f^{ij} = \frac{1}{3}((\frac{\tau^*}{2} + \tau) \delta_k^j + \tau^* S_k^j). \tag{29}$$

After substituting (29) in (27), we get

$$R_k^h = \frac{\tau}{3} \delta_k^h + \frac{\tau^*}{6} \Phi_k^h,$$

and also

$$R_{ki} = \frac{\tau}{3} g_{ki} + \frac{\tau^*}{6} f_{ki}, \quad R_{ki} f^{ih} = \frac{\tau}{6} S_k^h + \frac{\tau^*}{6} \delta_k^h. \tag{30}$$

From the last equation we find that

$$\tau^* = -\tau.$$

The equations (30) get the form

$$R_{ki} = \frac{\tau}{6}(2g_{ki} - f_{ki}), \quad R_{ki}f^{ih} = \frac{\tau}{6}(S_k^h - \delta_k^h). \tag{31}$$

Finely we obtain:

$$R_{ijk}^h = \frac{\tau}{6}(2\delta_k^h g_{ij} - 2\delta_j^h g_{ki} + (\delta_k^h + S_k^h)f_{ij} - (\delta_j^h + S_j^h)f_{ki})$$

and

$$R_{hijk} = \frac{\tau}{6}(2g_{kh}g_{ij} - 2g_{hj}g_{ki} + f_{kh}f_{ij} - f_{hj}f_{ki}).$$

The last identity is equivalent to (21).

We note that $R_{ijk}^h \neq 0$, so ∇ isn't a locally flat connection. □

Let p be a point in M and x, y be two linearly independent vectors in T_pM. It is known the value

$$\mu(x,y) = \frac{R(x,y,x,y)}{g(x,x)g(y,y) - g^2(x,y)}$$

is the sectional curvature of 2-section $\{x,y\}$.

Corollary 2.3. *Let M satisfy the conditions of the Theorem 2.2. Let x be an arbitrary vector in T_pM, and φ be the angle between x and qx. Then the sectional curvature of a 2-section $\{x,qx\}$ is*

$$\mu(x,qx) = -\frac{\tau}{6}\tan^2\frac{\varphi}{2}, \qquad \varphi \in (0, \frac{2\pi}{3}).$$

Corollary 2.4. *Let M satisfy the conditions of the Theorem 2.2. Then the Ricci tensor of g is degenerated.*

The proof follows from (31).

Note. Let $\{x,qx\}$ be a 2-section and $g(x,qx) = 0$. Then

$$\mu(x,qx) = -\frac{\tau}{6}.$$

REFERENCES

[1] G. Dzhelepov, I. Dokuzova, D. Razpopov. *On a three dimensional Riemannian manifold with an additional structure.* arXiv:math.DG/0905.0801

[2] G. Dzhelepov, D. Razpopov I. Dokuzova. *Almost conformal transformation on Riemannian manifold with an additional structure.* arXiv:math.DG/1010.4975

[3] K. Yano. *Differential geometry.* New York, Pergamont press, 1965

Iva Dokuzova
University of Plovdiv
FMI, Department of Geometry
236 Bulgaria Blvd.
Bulgaria 4003

e-mail:dokuzova@uni-plovdiv.bg

Dimitar Razpopov
Department of Mathematics and Physics
Agricultural University of Plovdiv
12 Mendeleev Blvd.
Bulgaria 4000
e-mail:drazpopov@qustyle.bg

Mathematics Subject Classification (2010): 53C15, 53B20

Keywords: Riemannian manifold, affinor structure, curvatures